

1-1-2010

Q fever cholecystitis in an unvaccinated butcher diagnosed by gallbladder polymerase chain reaction

Mel Figtree
St. George Hospital

Spiros Miyakis
St. George Hospital, smiyakis@uow.edu.au

John Stenos
Australian Rickettsial Reference Laboratory, Geelong

Stephen Graves
Australian Rickettsial Reference Laboratory, Geelong

Susan Botham
South Eastern Sydney And Illawarra Health

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/medpapers>



Part of the [Medicine and Health Sciences Commons](#)

Citation

Figtree, Mel; Miyakis, Spiros; Stenos, John; Graves, Stephen; Botham, Susan; Ferson, Mark; and Krilis, Steven, 2010, Q fever cholecystitis in an unvaccinated butcher diagnosed by gallbladder polymerase chain reaction, 421-423.
<https://ro.uow.edu.au/medpapers/562>

Q fever cholecystitis in an unvaccinated butcher diagnosed by gallbladder polymerase chain reaction

Abstract

Acalculous cholecystitis has been described in association with Q fever. We describe the first known case of Q fever cholecystitis diagnosed by polymerase chain reaction on gallbladder tissue. The presence of *Coxiella burnetii* genome in the diseased gallbladder tissue argues for direct involvement of the organism in the pathogenesis of Q fever cholecystitis.

Keywords

Q fever, Polymerase chain reaction, Cholecystitis

Disciplines

Medicine and Health Sciences

Publication Details

Figtree, M., Miyakis, S., Stenos, J., Graves, S., Botham, S., Ferson, M. & Krilis, S. A. (2010). Q fever cholecystitis in an unvaccinated butcher diagnosed by gallbladder polymerase chain reaction. *Vector Borne and Zoonotic Diseases*, 10 (4), 421-423.

Authors

Mel Figtree, Spiros Miyakis, John Stenos, Stephen Graves, Susan Botham, Mark Ferson, and Steven Krilis

Q Fever Cholecystitis in an Unvaccinated Butcher Diagnosed by Gallbladder Polymerase Chain Reaction

Mel Figtree,¹ Spiros Miyakis,¹ John Stenos,² Stephen Graves,² Susan Botham,³ Mark Ferson,³ and Steven Krillis¹

Abstract

Acalculous cholecystitis has been described in association with Q fever. We describe the first known case of Q fever cholecystitis diagnosed by polymerase chain reaction on gallbladder tissue. The presence of *Coxiella burnetii* genome in the diseased gallbladder tissue argues for direct involvement of the organism in the pathogenesis of Q fever cholecystitis.

Key Words: Q fever—Polymerase chain reaction—Cholecystitis.

THIS IS THE FIRST KNOWN REPORT of Q fever cholecystitis diagnosed by polymerase chain reaction (PCR) on the surgically removed gall bladder. A 38-year-old man was admitted to a Sydney teaching hospital with a 6-day history of fever, lethargy, myalgia, headache, and vomiting. He had worked for 6 months as a meat packer at a wholesale butcher. Workplace investigation revealed that his job entailed cutting frozen carcasses of cattle, sheep, and pigs with a high-speed saw, a process that generates spray, despite which no personal precautions against aerosol exposure were utilized. There was no history of Q fever vaccination. On examination the patient was febrile to 39.9°C, and had severe headache with mild photophobia and nonspecific abdominal tenderness.

Transaminases were raised with alanine aminotransferase 148 U/L (<45 U/L) and aspartate aminotransferase 150 U/L (<45 U/L). Alkaline phosphatase, gammaglutamyl transferase, and bilirubin were normal. C reactive protein was raised at 76 mg/L (<3 mg/L), and leucocyte count was 6.23×10^9 /L (3.50 – 11.00×10^9 /L) with a lymphopenia of 0.8×10^9 /L (1.5 – 4.0×10^9 /L). Cultures from blood, urine, and cerebrospinal fluid (CSF) were sterile. A presumptive diagnosis of Q fever was made on day 3, and the patient was commenced on doxycycline.

The patient's condition initially improved. However, on day 6 he developed severe right upper quadrant abdominal pain and fever to 38.5°C. Transaminases increased (alanine aminotransferase 325 U/L and aspartate aminotransferase

349 U/L), with evidence of cholestasis: bilirubin 61 μ mol/L (0–25 μ mol/L), alkaline phosphatase 287 U/L (38–126 U/L), and gammaglutamyl transferase 222 U/L (0–50 U/L). Abdominal ultrasonography revealed hepatomegaly and acalculous cholecystitis with a thickened edematous gallbladder measuring 11 mm (see Fig. 1). Intravenous ampicillin, gentamicin, and metronidazole were commenced, and the patient underwent urgent laparoscopic cholecystectomy given concern for possible gangrenous gallbladder. Histopathology showed acute cholecystitis with a diffuse infiltrate of lymphocytes, plasma cells, and occasional neutrophils. The patient defervesced postoperatively and was discharged 5 days after operation. Q fever was diagnosed by positive PCR from gallbladder tissue. Initial screening PCR assay using a *Com 1*-gene target was positive, and this was confirmed with a second PCR assay targeting the IS1111, a sequence in the *Coxiella burnetii* genome. Serological testing showed a rise in complement fixation titer from <8 to >256 after 11 days. PCR on blood from admission and day 6 were subsequently tested, and *C. burnetii* genome was detected on both assays.

Q fever is a zoonosis caused by *C. burnetii*. Q fever has a worldwide distribution, with the exception of New Zealand. The majority of human infections are acquired by inhalation of small particle droplets from urine, feces, milk, or bodily fluid from cattle, sheep, goats, or native animals. *C. burnetii* causes an initial bacteremia with widespread tissue seeding and consequently has a diverse range of clinical manifestations.

¹Department of Immunology, Allergy, and Infectious Disease, St. George Hospital, Sydney, Australia.

²Australian Rickettsial Reference Laboratory, Geelong, Australia.

³Department of Public Health Unit, South Eastern Sydney and Illawarra Health, Sydney, Australia.




FIG. 1. Abdominal ultrasonography of patient showing acalculous cholecystitis. Thickened edematous gallbladder wall measuring 11 mm.

Acalculous cholecystitis has been described in association with Q fever (Rolain et al. 2003). Acalculous cholecystitis is a condition most commonly seen in patients critically unwell from a variety of conditions, including sepsis, trauma, or burns. Acalculous cholecystitis has been described as a direct complication of several microorganisms, including *Salmonella* spp., *Mycobacterium* spp., *Plasmodium* spp., *Candida* spp., cytomegalovirus, cryptosporidium, and microspora (Barie and Eachempati 2003). Such cases have been successfully managed with medical therapy directed at the specific organism.

Fifteen cases of Q fever-associated cholecystitis have been described (Kelly et al. 1986, Isaacs and Donald 1998, Modol et al. 1999, Rolain et al. 2003, Gonzalez-Delgado et al. 2005, Reina-Serrano et al. 2005, Hartzell et al. 2007). Two cases were diagnosed by liver biopsy with histopathology showing classic doughnut hole and fibrin ring surrounding a granuloma. The remainder were diagnosed by serology weeks after the initial illness. Immunohistochemistry was nondiagnostic for Q fever in all six patients who underwent cholecystectomy.

This is the first case described of Q fever cholecystitis diagnosed from gallbladder tissue. The presence of *C. burnetii* genome in this patient, together with the clinical presentation, acute cholecystitis, and seroconversion, suggests that *C. burnetii* may be directly involved in the pathogenesis of cholecystitis.

Molecular techniques are evolving as useful tools to make an early diagnosis of Q fever. New techniques utilizing htpAB-associated repetitive element of *C. burnetii* genome (IS1111) have been shown in an outbreak setting to have a sensitivity of 59% on serum (Turra et al. 2006). Sensitivity is highest in early infection, before immunoglobulin M rise. Molecular techniques may allow a more rapid diagnosis of Q fever cholecystitis, facilitating earlier implementation of specific treatment and reducing the likelihood of operation. Despite early medical management with doxycycline, patients

may still develop complications such as secondary bacterial infection, abscess formation, and perforation necessitating surgical intervention.

The annual incidence of Q fever cases notified in Australia has decreased from 5.0 per 100,000 in 1993, to 2.2 cases per 100,000 persons in 2004 (Brotherton et al. 2007). This reduction has been attributed to implementation between 2001 and 2004 of the National Q Fever Management Program, which targeted abattoir workers, those contracted to abattoirs, sheep shearers, and sheep, dairy, and cattle farmers and their employees. The patient described in this case would not have been identified as part of the program but was in a high-risk occupational group for which immunization is recommended. Australian guidelines recommend Q fever vaccine for abattoir workers, farmers, stockyard workers, shearers, animal transporters, others exposed to cattle, sheep, goats, and kangaroos or their products, and veterinarians and laboratory staff handling *C. burnetii* specimens. In this instance, a recommendation was made to the employer by Work Cover New South Wales that employees working in the meat-packing industry with negative Q fever serology be vaccinated and personal protective equipment (goggles, mask, gown, and gloves) be worn when operating the band saw.

Ongoing vigilance is required to prevent Q fever in high-risk groups. Of concern is the high proportion of Q fever cases occurring in groups with no recognized risk factors. Of the 15 described cases of Q fever-associated cholecystitis, only 4 had documented animal exposure. The proportion of Australian Q fever cases in people with no direct animal contact may increase during drought, as the subsequent dusty conditions enhance dissemination of this organism.

Acknowledgment

Phill Cantrell, Senior Project Officer, Hazard Management Group, WorkCover, New South Wales, Australia.

Disclosure Statement

No competing financial interests exist.

References

- Barie, PS, Eachempati, SR. Acute acalculous cholecystitis. *Curr Gastroenterol Rep* 2003; 5:302–309.
- Brotherton, J, Wang, H, Schaffer, A, Quinn, H, et al. Vaccine preventable diseases and vaccination coverage in Australia, 2003 to 2005. *CDI* 2007; 31(Suppl):S1–S152.
- Gonzalez-Delgado, L, Lopez, LG, Santolaria, PS, Garcia, PD, et al. Acalculous cholecystitis: an uncommon form of presentation of Q fever. *Gastroenterol Hepatol* 2005; 28:232–236.
- Hartzell, JD, Peng, SW, Wood-Morris, RN, Sarmiento, DM, et al. Atypical Q fever in US soldiers. *Emerg Infect Dis* 2007; 13:1247–1249.
- Isaacs, D, Donald, P. Brothers with fever and abdominal pain. *Pediatr Infect Dis J* 1998; 17:665–666.
- Kelly, R, Byrnes, D, Turner, J. Acute, severe hepatitis due to *Coxiella burnetii* infection. *Med J Aust* 1986; 144:151–155.
- Modol, JM, Llamazares, JF, Mate, JL, Troya, J, et al. Acute abdominal pain and Q fever. *Eur J Clin Microbiol Infect Dis* 1999; 18:158–160.

- Reina-Serrano, S, Jimenez-Saenz, M, Herrerias-Gutierrez, JM, Venero-Gomez, J. Q fever-related cholecystitis: a missed entity? *Lancet Infect Dis* 2005; 5:734–735.
- Rolain, JM, Lepidi, H, Harle, JR, Allegre, T, et al. Acute acalculous cholecystitis associated with Q fever: report of seven cases and review of the literature. *Eur J Clin Microbiol Infect Dis* 2003; 22:222–227.
- Turra, M, Chang, G, Whybrow, D, Higgins, G, et al. Diagnosis of acute Q fever by PCR on sera during a recent outbreak in rural south Australia. *Ann NY Acad Sci* 2006; 1078:566–569.

Address correspondence to:

Mel Figtree

Department of Immunology, Allergy, and Infectious Disease

St. George Hospital

1st Floor, 2 South Street

Kogarah

Sydney 2217

New South Wales

Australia

E-mail: melfigtree@yahoo.com.au

